



Paper Type: Original Article

## Neutrosophic Sober Open Sets: An Overview

Subasree Ramachandran<sup>1,\*</sup> , BasariKodi K<sup>1</sup>, Saeid Jafari<sup>2</sup>, Subramanian K<sup>1</sup>, Poornima R<sup>3</sup>, Alapati Varalakshmi<sup>4</sup>

<sup>1</sup>Department of Mathematics, Ramco Institute of Technology, Rajapalayam, India; subasree@ritrjpm.ac.in; basarikodi@ritrjpm.ac.in; subramanian@ritrjpm.ac.in.

<sup>2</sup>College of Vestsjaelland South & Mathematical and Physical Science Foundation, 4200 Slagelse, Denmark; jafaripersia@gmail.com.

<sup>3</sup>Department of Science and Humanities, Hindusthan College of Engineering and Technology, Coimbatore, India; poornimavisaanth@gmail.com.

<sup>4</sup>Manipal Academy of Higher Education, Manipal, India; lakshmi.alapati@manipal.edu.

### Citation:

Received: 26 November 2024	Ramachandran, S., BasariKodi, K., Jafari, S., Subramanian, K., Poornima, R., & Varalakshmi, A. (2025). Neutrosophic sober open sets: An overview. <i>Uncertainty discourse and applications</i> , 2(2), 139-145.
Revised: 27 February 2025	
Accepted: 28 May 2025	

### Abstract

In this study, a new class of neutrosophic sets, known as Neutrosophic sober open sets in Neutrosophic topological spaces, is introduced, and its fundamental features are examined. After this, we create a new topology type, known as a neutrosophic sober topology, in order to study its relationships in more detail. The introduced neutrosophic sober open sets and the associated sober topology provide a structured framework that enriches neutrosophic topological spaces by refining openness through relational constraints. The developed interior and closure operators further confirm the internal consistency of this framework and its capability to capture nuanced topological behavior under indeterminacy. These results highlight the potential of neutrosophic sober structures for advancing generalized topology under uncertainty.

**Keywords:** Neutrosophic set, Neutrosophic sober open set, Neutrosophic sober topological space,  $N\check{S}(Cl)$  operator,  $N\check{S}(Int)$  operator.

## 1 | Introduction

Zadeh [1] introduced fuzzy sets in 1965, which made it possible for things to belong to the unit interval [0,1] to varying degrees of membership. Atanassov [2] expanded this idea in 1983 by proposing Intuitionistic Fuzzy Sets (IFS) [3], which take into account both degrees of membership and non-membership. In 1998, Smarandache [4] presented neutrosophic sets, which were created by adding indeterminacy as a component to the truth and falsity membership functions [5].

 Corresponding Author: subasree@ritrjpm.ac.in

 <https://doi.org/10.48313/uda.vi.65>

 Licensee System Analytics. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>).

Several writers have recently proposed new forms of sets in general topology [6]. A novel class of sets, known as sober open sets in sober topological spaces, was recently introduced by Jayanthi and Jafari [7] in 2024. Additionally, counterexamples have been used to illustrate the sober continuity, sober connectedness, and separation axioms.

This encourages us to explore the basic features of sober open sets in neutrosophic topological spaces. These concepts have been illustrated with an example. Furthermore, the closure and interior operators of neutrosophic sober topological spaces have been introduced and studied.

## 2 | Preliminaries

**Definition 1 ([4]).** Let  $X$  be a fixed, non-empty set. A set with the form  $N = \{\langle x, T_N(x), I_N(x), F_N(x) \rangle : x \in X\}$  is called a Neutrosophic set, where  $T_N(x)$ ,  $I_N(x)$ ,  $F_N(x)$  represent the degree of truth, degree of indeterminacy and the degree of falsity respectively of each element  $x \in X$  to the set  $N$ .

**Definition 2 ([8]).** The complement of a Neutrosophic set  $N$  is denoted by  $N^c$  and is defined by  $N^c = \{\langle x, F_A(x), 1-I_A(x), T_A(x) \rangle : x \in X\}$ .

**Definition 3 ([8]).** Consider two Neutrosophic sets  $U$  and  $V$  over  $X$ , then  $U$  is said to be contained in  $V$ , denoted by  $U \subseteq V$ , if and only if  $T_u(x) \leq T_v(x)$ ,  $I_u(x) \leq I_v(x)$ ,  $F_u(x) \geq F_v(x)$ .

**Definition 4 ([8]).** The arbitrary union of two Neutrosophic sets  $U$  and  $V$  over  $X$  is denoted by  $U \cup V$  and is defined by  $\{\langle x, T_u(x) \vee T_v(x), I_u(x) \vee I_v(x), F_u(x) \wedge F_v(x) \rangle : x \in X\}$ .

**Definition 5 ([8]).** The finite intersection of two Neutrosophic sets  $U$  and  $V$  over  $X$  is denoted by  $U \cap V$  and is defined by  $\{\langle x, T_u(x) \wedge T_v(x), I_u(x) \wedge I_v(x), F_u(x) \vee F_v(x) \rangle : x \in X\}$ .

**Definition 6 ([8]).** Let  $N$  be a Neutrosophic set over  $X$ , then the universe set of  $N$  is denoted by  $1_N$  and is defined by  $1_N = \{\langle x, 1, 1, 0 \rangle : x \in X\}$ .

**Definition 7 ([8]).** Let  $N$  be a Neutrosophic set over  $X$ , then the empty set of  $N$  is denoted by  $0_N$  and is defined by  $0_N = \{\langle x, 0, 0, 1 \rangle : x \in X\}$ .

**Definition 8 ([8]).** Let  $U$  and  $V$  be two Neutrosophic sets over  $X$ , the difference of  $U$  and  $V$  is denoted by  $D_{uv} = U \setminus V$  and is defined by  $\{\langle x, T_u(x) \wedge F_v(x), I_u(x) \wedge (1-I_v(x)), F_u(x) \vee T_v(x) \rangle : x \in X\}$ .

**Definition 9 ([9]).** A Neutrosophic Topology (NT) on a non-empty set  $X$  is a family  $\tau_N$  of neutrosophic subsets in  $X$  that satisfies the following axioms [10]:

- I.  $0_N, 1_N \in \tau_N$ .
- II.  $U \cap V \in \tau$  for any  $U, V \in \tau_N$ .
- III.  $\bigcup N_i \in \tau$  for all  $\{N_i : i \in J\} \subseteq \tau_N$ .

The pair  $(X, \tau_N)$  represents a neutrosophic topological space  $\tau_N$  over  $X$ .

We denote a neutrosophic open set as an NO set and a neutrosophic topological space as NTS.

## 3 | Neutrosophic Sober Open Sets

The definitions of a new open set in Neutrosophic topological space, known as a Neutrosophic sober open set and its complement, a Neutrosophic sober closed set, are presented in this section along with a discussion of their fundamental characteristics. To further investigate Neutrosophic sober open sets, a Neutrosophic topological space known as the Neutrosophic sober topological space is also introduced.

**Definition 10.** A non-empty set  $N \neq X$  is said to be a Neutrosophic sober open set in Neutrosophic topological space  $(X, \tau_N)$ , if there exist two distinct non-empty NO sets  $N_1 \neq X$  and  $N_2 \neq X$  in  $X$  that satisfy the conditions:

- I.  $N \cup N1$  is neutrosophic open and  $N \cup N1 \neq X$ .
- II.  $N \cap N2$  is neutrosophic open and  $N \cap N2 \neq \phi$ .

The complement  $N^c$  of a neutrosophic sober open set  $N$  is a neutrosophic sober closed set in  $(X, \tau_N)$ . The collection of all neutrosophic sober open (resp. neutrosophic sober closed) sets is denoted by  $NS-O(X)$  (resp.  $NS-C(X)$ ).

**Example 1.** Let  $X = \{a, b\}$  and  $\tau_N = \{0_N, A, B, C, D, 1_N\}$  be a neutrosophic topological space, where

$A = \langle (a, 0.3, 0.3, 0.5), (b, 0.4, 0.4, 0.6) \rangle$ ,  $B = \langle (a, 0.1, 0.2, 0.3), (b, 0.4, 0.5, 0.6) \rangle$ ,  $C = \langle (a, 0.1, 0.2, 0.5), (b, 0.4, 0.4, 0.6) \rangle$  and  $D = \langle (a, 0.3, 0.3, 0.3), (b, 0.4, 0.5, 0.6) \rangle$ .

Here,  $P = \langle (a, 0.3, 0.2, 0.5), (b, 0.4, 0.4, 0.6) \rangle$  is an NS-open set since there exist two non-empty NO sets  $A$  and  $B$  such that  $P \cup A = A \neq X$  and  $P \cap B = C \neq \phi$ .


$Q = \langle (a, 0.1, 0.3, 0.3), (b, 0.4, 0.5, 0.6) \rangle$  is an NS-open set since there exist two non-empty NO sets  $D$  and  $B$  such that  $Q \cup D = D \neq X$  and  $Q \cap B = B \neq \phi$ .

$R = \langle (a, 0.3, 0.3, 0.5), (b, 0.4, 0.5, 0.6) \rangle$  is an NS-open set since there exist two non-empty NO sets  $D$  and  $A$  such that  $R \cup D = D \neq X$  and  $R \cap A = A \neq \phi$ .

$S = \langle (a, 0.2, 0.2, 0.3), (b, 0.4, 0.5, 0.6) \rangle$  is an NS-open set since there exist two non-empty NO sets  $A$  and  $C$  such that  $S \cup A = D \neq X$  and  $S \cap C = C \neq \phi$ .

It is comprehensible that NS-open sets  $P, Q, R,$  and  $S$  are such that  $C \subseteq P \subseteq A, B \subseteq Q \subseteq D, A \subseteq R \subseteq D,$  and  $C \subseteq S \subseteq D$ .

**Remark 1.** A Hasse diagram is one way to represent the NS-open sets in *Example 1* with the relation  $\subseteq$ .



**Fig. 1.** Hasse diagram of neutrosophic sober open sets with respect to inclusion.

**Definition 11.** The union of all neutrosophic sober open sets (resp. neutrosophic sober closed sets) in  $X$  is denoted by  $\bigcup \{NS-O(X)\}$  (resp.  $\bigcup \{NS-C(X)\}$ ).

**Definition 12.**

- I. The neutrosophic sober complement of NS-O set  $U$  is defined by  $(NS)^c(U) = \bigcup \{NS-O(X)\} \setminus U$ .
- II. The neutrosophic sober complement of NS-C set  $V$  is defined by  $(NS)^c(V) = \bigcup \{NS-C(X)\} \setminus V$ .

**Example 2.** This example shows that the neutrosophic sober complement of NS-O set need not be neutrosophic sober closed.

From *Example 1*, the union of all NS-open sets is  $D$ .

$(NS)c(P) = D \setminus P = D$  which is not neutrosophic sober closed.

**Remark 2.** The neutrosophic sober complement of the NS-C set is always neutrosophic sober open.

From *Example 1*, the union of all NS-closed sets is  $C_c$ .

$(NS)c(P_c) = C_c \setminus P_c = P$  which is neutrosophic sober open.

$(NS)c(Q_c) = C_c \setminus Q_c = Q$  which is neutrosophic sober open.

$(NS)c(R_c) = C_c \setminus R_c = R$  which is neutrosophic sober open.

$(NS)c(S_c) = C_c \setminus S_c = S$  which is neutrosophic sober open.

**Example 3.** Every NO set in  $(X, \tau_N)$  need not be an NS-O set in general.

Let  $X = \{u, v, w\}$  and  $\tau_N = \{0N, N, 1N\}$  be a neutrosophic topological space, where  $N = \langle (u, 0.5, 0.6, 0.4), (v, 0.4, 0.5, 0.2), (w, 0.7, 0.6, 0.9) \rangle$ .

Here  $N$  is a NO set but not an NS-O set, since there does not exist two non-empty disjoint NO sets such that  $N \cup N_1 \neq X$  and  $N \cap N_2 \neq \phi$ .

**Theorem 1.** Necessary and sufficient condition for a NO set to be an NS-O set.

A NO set  $N$  in  $(X, \tau_N)$  is NS-open if and only if there exists a proper NO set  $M$  such that  $N \subseteq M$  or  $M \subseteq N$ .

Proof:

**Necessity.** Let  $N$  be a neutrosophic open set in  $(X, \tau_N)$  that is NS-open, then by *Definition 10*, there exist two proper NO sets, say  $N$  itself and  $M$ , such that  $N \subseteq N \subseteq M$  or  $M \subseteq N \subseteq N$ .

**Sufficiency.** Assume that  $N$  is a NO set in  $(X, \tau_N)$  and that if  $N \subseteq M$ , where  $M$  is a proper NO set in  $X$ . Then we have the conditions  $N \cup N = N \neq X$  and  $N \cap M = N \neq \phi$ , both of which are NO in  $X$ . If on the other hand,  $M \subseteq N$ , then we have  $N \cup N = N \neq X$  and  $N \cap M = M \neq \phi$ , both of which are NO in  $X$ . In both cases,  $N$  is NS-open in  $X$ .

**Remark 3.** It is obvious that  $1N$  and  $0N$  are not neutrosophic sober open sets.

**Theorem 2.** A chain has an NS-open set if it contains at least two non-disjoint NO sets.

Proof:



We know that in a chain, every element is comparable. That is, we consider an NTS with two non-disjoint NO sets, say  $A$  and  $B$ , which can be comparable, then there exists a NS-open set  $U$  such that  $A \subseteq U \subseteq B$ . It is clearly shown in the following figure.

**Fig. 2. Existence of a neutrosophic sober open set  $U$  in a chain of neutrosophic open sets between two non-disjoint sets  $A$  and  $B$ .**

**Remark 4.** It is evident that a Neutrosophic topological space  $(X, \tau_N)$  with a single NO set does not contain any neutrosophic sober open sets.

### 4 | Neutrosophic Sober Topological Space

**Definition 13.** We shall now consider an NTS  $(X, \tau_N)$  with at least two non-disjoint NO sets. In this NTS, the following conditions hold:

- I.  $0_N, 1_N \in \tau_N$ .
- II. The arbitrary union of two NS-open sets is again NS-open in  $(X, \tau_N)$ .
- III. The finite intersection of two NS-open sets is again NS-open in  $(X, \tau_N)$ .

We refer to the above NTS as a Neutrosophic sober topological space, and it is denoted as  $\tau_{NSober}$  and the corresponding space as  $X_{NSober}$ .

**Definition 14.** Let  $(X_{NSober}, \tau_{NSober})$  be a Neutrosophic sober topological space. Then the Neutrosophic sober interior and Neutrosophic sober closure of any proper set  $A$  in  $(X_{NSober}, \tau_{NSober})$  are denoted by  $N\check{S}(Int)$  and  $N\check{S}(Cl)$ , respectively, and are defined by


$$N\check{S} - Int (A) = \bigcup \{G : G \text{ is NS-open in } X_{NSober} \text{ and } G \subseteq A\}.$$

$$N\check{S} - Cl (A) = \bigcap \{K : K \text{ is NS-closed in } X_{NSober} \text{ and } A \subseteq K\}.$$

**Remark 5.** For any proper set  $A$ ,  $N\check{S} - Int (A) \subseteq A$  is true only when  $N\check{S} - Int (A)$  exists. Also, if  $A \subseteq B \Rightarrow N\check{S} - Int (A) \subseteq N\check{S} - Int (B)$  is true for any two proper sets  $A$  and  $B$ , only when  $N\check{S} - Int (A)$  and  $N\check{S} - Int (B)$  exist.

**Example 4.** Let  $X = \{a, b\}$  and  $\tau_{NSober} = \{0_N, A, B, C, D, 1_N\}$  be a neutrosophic topological space, where  $A = \langle (a, 0.5, 0.5, 0.3), (b, 0.3, 0.3, 0.2) \rangle$ ,  $B = \langle (a, 0.2, 0.2, 0.1), (b, 0.7, 0.7, 0.6) \rangle$ ,  $C = \langle (a, 0.2, 0.2, 0.3), (b, 0.3, 0.3, 0.6) \rangle$  and  $D = \langle (a, 0.5, 0.5, 0.1), (b, 0.7, 0.7, 0.2) \rangle$ .

Here  $P = \langle (a, 0.4, 0.4, 0.3), (b, 0.3, 0.3, 0.4) \rangle$ ,  $Q = \langle (a, 0.2, 0.2, 0.2), (b, 0.5, 0.5, 0.6) \rangle$ ,  $R = \langle (a, 0.3, 0.3, 0.1), (b, 0.7, 0.7, 0.5) \rangle$ ,  $S = \langle (a, 0.5, 0.5, 0.2), (b, 0.6, 0.6, 0.2) \rangle$  and  $T = \langle (a, 0.4, 0.4, 0.2), (b, 0.5, 0.5, 0.5) \rangle$  are some of the NS-open sets.



**Fig. 3.** Hasse diagram representation of the neutrosophic sober topology  $\tau_{NSober}$  on the set  $X = \{a, b\}$ , illustrating the NS-open sets  $A, B, C, D, P, Q, R, S$ , and  $T$ .

Now we find the  $N\check{S} - Int (A)$  for any proper set of  $\tau_{NSober}$ .

**Table 1. Neutrosophic sober interior of selected NS-open sets and proper sets in the neutrosophic sober topological space  $(X, \tau_{NSoBer})$ .**

NS-Open Sets	$N\check{S} - Int (NS\text{-Open Set})$	Proper Set	$N\check{S} - Int (Proper Set)$
P	P	A	P
Q	Q	B	Q
R	R	C	DOES NOT EXIST
S	S	D	D
T	T		

**Theorem 3.** Let  $(X_{NSoBer}, \tau_{NSoBer})$  be a Neutrosophic sober topological space. Then for any proper set in  $X_{NSoBer}$ , the following holds:

- I.  $N\check{S} - Int (A) \subseteq A$ .
- II.  $A \subseteq B$  then  $N\check{S} - Int (A) \subseteq N\check{S} - Int (B)$ .
- III.  $N\check{S} - Int (N\check{S} - Int (A)) = N\check{S} - Int (A)$ .
- IV.  $N\check{S} - Int (M \cap N) = N\check{S} - Int (M) \cap N\check{S} - Int (N)$ .
- V.  $N\check{S} - Int (M) \cup N\check{S} - Int (N) \subseteq N\check{S} - Int (M \cup N)$ .

Proof: The results are apparent from the above example.

## 5 | Conclusion

Neutrosophic sober open sets are introduced in this article, along with relevant examples that illustrate their fundamental properties. The concepts of neutrosophic sober interior and closure operators are further discussed in order to support the neutrosophic sober topology. The theorems and properties of neutrosophic sober-open sets in neutrosophic sober topology are explained with related examples. There is also an extent to incorporate continuous functions, connectedness, and compactness based on neutrosophic sober topological spaces.

## Acknowledgments

Acknowledgments enable you to thank all those who have contributed to the research. Careful thought needs to be given concerning those whose help should be acknowledged and in what order. The general advice is to express your appreciation concisely and to avoid strong emotive language.

## Authors' Contributions

S. R.: Research Design, Conceptualization, and Validation. B. K. K.: Data Curation, Computing, and Editing. S. J.: Methodology, Visualization and Formal Analysis. S. K.: Research Design and Validation P. R.: Visualization and Formal Analysis. A. V.: Methodology and Formal Analysis. The authors have read and agreed to the published version of the manuscript.

## Consent for Publication

The authors confirm consent for the publication of this work.

## Ethics Approval and Consent to Participate

The authors confirm that this research did not involve human participants or animal subjects.

## Funding

This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Data Availability

All data are included in the text.

## Conflict of Interest

The authors declare that they have no conflict of interest.

## References

- [1] Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8(3), 338–353. [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X)
- [2] Atanassov, K. T., & Stoeva, S. (1986). Intuitionistic fuzzy sets. *Fuzzy sets and systems*, 20(1), 87–96. [https://doi.org/10.1016/S0165-0114\(86\)80034-3](https://doi.org/10.1016/S0165-0114(86)80034-3)
- [3] Çoker, D. (1997). An introduction to intuitionistic fuzzy topological spaces. *Fuzzy sets and systems*, 88(1), 81–89. [https://doi.org/10.1016/S0165-0114\(96\)00076-0](https://doi.org/10.1016/S0165-0114(96)00076-0)
- [4] Smarandache, F. (1999). A unifying field in logics: Neutrosophic logic. In *Neutrosophy, neutrosophic set, neutrosophic probability* (pp. 1–141). American Research Press. <https://web-archive.southampton.ac.uk/cogprints.org/1919/3/eBook-Neutrosophics2.pdf>
- [5] Salama, A. A., Smarandache, F., & Kromov, V. (2014). Neutrosophic closed set and neutrosophic. *Neutrosophic sets and systems*, 4, 4–8. <https://fs.unm.edu/NeutrosophicClosedSet.pdf>
- [6] Al-Hamido, R. K. (2024). Neutrosophic generalized topological space. *Plithogenic logic and computation*, 2, 1–9. <https://orcid.org/0000-0001-8388-0557>
- [7] Jayanthi, D., & Jafari, S. (2024). Sober open sets in sober topological spaces. *South east asian journal of mathematics and mathematical sciences*, 20(3), 385–396. <https://doi.org/10.56827/SEAJMMS.2024.2003.29>
- [8] Wang, H., Smarandache, F., Zhang, Y., & Sunderraman, R. (2010). *Single valued neutrosophic sets*. Infinite Study. <https://fs.unm.edu/SingleValuedNeutrosophicSets.pdf>
- [9] Salama, A. A., & Alblowi, S. A. (2012). Neutrosophic set and neutrosophic topological spaces. *IOSR journal of mathematics*, 3(4), 31–35. <https://doi.org/10.9790/5728-0343135>
- [10] James R, M. (2004). *Topology*. Prentice –Hall of India. <https://archive.org/details/topology0002edmunk>